

## LTCC Geometry and Physics: Exercise Sheet 2

(Note: Unless otherwise stated, the Einstein summation convention is assumed throughout.)

1. Given a Riemannian manifold  $M$  with metric  $g$ , consider the action  $S = \int_{t_0}^{t_1} L dt$  with the purely kinetic Lagrangian

$$L = \frac{1}{2}g(\dot{\mathbf{x}}, \dot{\mathbf{x}}) = \frac{1}{2}g_{jk}\dot{x}^j\dot{x}^k, \quad (1)$$

where  $g = (g_{jk}(\mathbf{x}))$  is the metric given in terms of local coordinates  $x^j$  for a point  $\mathbf{x} \in M$  with tangent vector  $\dot{\mathbf{x}}$ ; the dot denotes differentiation w.r.t. the parameter  $t$  along a curve.

(i) Applying the principle of least action,  $\delta S = 0$ , calculate the Euler-Lagrange equations for this Lagrangian, and show that they can be written in the form

$$\ddot{x}^j + \Gamma_{kl}^j \dot{x}^k \dot{x}^l = 0, \quad (2)$$

where  $\Gamma_{kl}^j$  (the Christoffel symbols) should be found in terms of  $g$  and its derivatives.

(ii) Perform a Legendre transformation and hence reformulate the geodesic equations as an Hamiltonian system on  $T^*M$ .

2. Write down the Euclidean metric on  $\mathbb{R}^3$  in Cartesian coordinates, and the invariant volume form. Use this to calculate the invariant 2-form on the sphere  $S^2$ , and show that this is a symplectic form.

3.(i) For a pair of vector fields  $X, Y$  on a smooth manifold  $M$  of dimension  $d$ , define their commutator  $[X, Y]$  (or Lie bracket) by the commutator of the corresponding differential operators. Show that with this definition the vector fields form a Lie algebra.

(ii) Now suppose that  $d = 2n$  and  $(M, \omega)$  is a symplectic manifold. For Hamiltonian vector fields  $X_G, X_H$  corresponding to smooth functions  $G, H$ , show that the following formula holds:

$$[X_G, X_H] = -X_{\{G, H\}},$$

where  $\{, \}$  denotes the Poisson bracket. (Hint: Use Darboux's theorem.)

(iii) Prove that the Hamiltonian vector fields form a Lie subalgebra of the vector fields.

4.(i) Check that the formula

$$\{\pi_j, \pi_k\} = -\epsilon_{jkl}\pi_l$$

defines a Poisson bracket on  $\mathbb{R}^3$  with coordinates  $(\pi_1, \pi_2, \pi_3)$ , and show that (on linear functions) this is isomorphic to the Lie algebra  $\mathfrak{so}(3)$  of the rotation group in three dimensions.

(ii) Write down the Poisson tensor for the above bracket and calculate its rank. Show that  $C = \pi_1^2 + \pi_2^2 + \pi_3^2$  is a Casimir for this bracket, and describe the symplectic leaves.

(iii) Consider the Hamiltonian system on  $\mathbb{R}^3$  defined by the Hamiltonian

$$H = \frac{\pi_1^2}{2I_1} + \frac{\pi_2^2}{2I_2} + \frac{\pi_3^2}{2I_3}.$$

Write down the equations of motion, and show that they are equivalent to the Euler top (free motion of a rigid body about a fixed point) in terms of the angular momentum  $\boldsymbol{\pi}$  and angular velocity  $\boldsymbol{\omega}$ , where  $\boldsymbol{\pi} = I\boldsymbol{\omega}$  with the inertia tensor  $I = \text{diag}(I_1, I_2, I_3)$ . Can you describe the orbits?